arXiv:1004.5494v2 [math.AP] 10 Jun 2010

On the nodal line of the second eigenfunction of the Laplacian over some concave domains in $\mathbb{R}^2$

Donghui Yang
School of Mathematics and Statistics, Central China Normal University,
430079, Wuhan, People's Republic of China
donghyang@mail.ccnu.edu.cn

Abstract

In this paper we will prove the nodal line N of the second eigenfunction of the Laplacian over some simply connected concave domain Ω in $\mathbb{R}^2$ must intersect the boundary $\partial\Omega$ at exactly two points.

1 Introduction

An eigenfunction φ_i is meant to be a solution of Dirichlet's problem:

$$\begin{cases} \Delta\varphi_i + \lambda_i\varphi_i = 0 & \text{in } \Omega \\ \varphi_i = 0 & \text{on } \partial\Omega, \end{cases} \tag{1.1}$$

where $\Delta = \dfrac{\partial^2}{\partial x_1^2} + \dfrac{\partial^2}{\partial x_2^2}$ is the Laplacian, Ω is a bounded smooth domain in $\mathbb{R}^2$, λ_i is the ith eigenvalue with $\lambda_1 < \lambda_2 \leq \lambda_3 \leq \cdots$, and φ_i is the ith eigenfunction $(i = 1, 2, \cdots)$. It is well know that the first eigenfunction is positive in Ω, and all higher eigenfunctions must change sign. The nodal set of an eigenfunction φ_i is defined to be the closure of $\{x \in \Omega;\ \varphi_i(x) = 0\}$. The Courant nodal domain theorem [2] tells us that the nodal set of an ith eigenfunction φ_i divides the domain Ω into at most i subregions. especially, φ_2 divides the domain Ω into at exactly 2 domains.

In 1967 Payne [11] conjectured that φ_2 cannot have a closed nodal line in Ω and in 1982 Yau [12] asked the same question for convex domains in $\mathbb{R}^2$. Payne [10] proved that the nodal line touches the boundary of a convex set which is symmetric under

---

[1] The authors were supported by NSFC 10901069.

a reflection. C.-S. Lin [8] proved the conjecture provided the domain $\Omega \subset \mathbb{R}^2$ is smooth, convex, and invariant under a rotation with angle $\frac{2\pi p}{q}$, where p and q are positive integers. D. Jerison [6] proved the conjecture for long thin convex sets. Melas [9] have settled the convex case for C^∞ boundary and this was extended to general boundary by Alessandrini [1]. M. Hoffmann-Ostenhof, T. M. Hoffmann-Ostenhof and N. Nadirashvili [5] construct a nonconvex, not simply connected domain for which the second eigenfunction has a closed nodal line. Also for convex D D. Jerison [7] and D. Grieser and D. Jerison [4] obtained interesting results on the location of the first nodal line.

In this paper we obtain that the nodal line of the second eigenfunction φ_2 over some simply connected concave domains Ω intersect the boundary $\partial\Omega$ at exactly two points. Which is the special case of the following theorem:

Theorem The nodal line of a second eigenfunction of Laplacian divides the domain Ω by intersecting its boundary at exactly two points if the domain $\rho(\Omega)$ is strictly convex in θ and symmetric with respect to the r-axis.

2 main results and their proofs

Let $\Omega, \widetilde{\Omega} \subset \mathbb{R}^2$ be two smooth domains (0 is not in the closure of $\widetilde{\Omega}$) and $\rho : \Omega \to \widetilde{\Omega}$, $(x, y) \mapsto (r, \theta)$ be a diffeomorphism defined by $(x, y) = \rho^{-1}(r, \theta)$ with $x = r\cos\theta, y = r\sin\theta$. Then the equation (1.1) becomes a new equation:

$$\begin{cases} \dfrac{\partial^2 \widetilde{\varphi}_i}{\partial r^2} + \dfrac{1}{r^2}\dfrac{\partial^2 \widetilde{\varphi}_i}{\partial \theta^2} + \dfrac{1}{r}\dfrac{\partial \widetilde{\varphi}_i}{\partial r} + \lambda_i \widetilde{\varphi}_i = 0 & \text{in } \widetilde{\Omega} \\ \widetilde{\varphi}_i = 0 & \text{on } \partial\widetilde{\Omega}, \end{cases} \tag{2.1}$$

where $\widetilde{\varphi}_i(r, \theta) = \varphi_i \circ \rho^{-1}(r, \theta)$.

Now we display the relations between the equations (1.1) and (2.1): If φ_i is a solution of equation (1.1), then φ_i is a smooth function and $\widetilde{\varphi}_i = \varphi_i \circ \rho^{-1}$ is also a smooth solution of (2.1); conversely, if $\widetilde{\varphi}_i$ is a solution of (2.1), then $\widetilde{\varphi}_i \in H_0^1(\widetilde{\Omega})$, and $\widetilde{\varphi}_i \in C^\infty(\widetilde{\Omega})$ by the infinite differentiability up to the boundary theorem (See [3] pp. 324-326), hence $\varphi_i = \widetilde{\varphi}_i \circ \rho$ is a solution of equation (1.1). And hence $\lambda_2(\Omega) = \lambda_2(\rho(\Omega))$.

Following from Courant nodal domain theorem [2] we know that the nodal set of an ith eigenfunction $\widetilde{\varphi}_i$ divides the domain $\widetilde{\Omega}$ into at most i subregions, especially, $\widetilde{\varphi}_2$ divides the domain $\widetilde{\Omega}$ into at exactly 2 subregions.

Throughout the paper we denote φ_2 an second eigenfunction of (1.1) and $N = \overline{\{x \in \Omega;\ \varphi_2(x) = 0\}}$ is the nodal line of φ_2; $\widetilde{\varphi}_2 = \varphi_2 \circ \rho^{-1}$ and $\widetilde{N} = \overline{\{x \in \widetilde{\Omega};\ \widetilde{\varphi}_2(x) = 0\}}$.

The following Lemma 1 is proved in [8]:

Lemma 1 Suppose $P \in \partial\Omega$. Then $\dfrac{\partial \varphi_2}{\partial \nu}(P) = 0$ if and only if $P \in N$, where $\dfrac{\partial \varphi_2}{\partial \nu}$ is the outnormal derivative of φ_2 on the boundary.

Proof Now, we prove Lemma 1 again as some different way.

Let $P \in \partial\Omega$. Since φ_2 is a smooth solution of equation (1.1) and Ω is a smoothly bounded domain, there exists an open set $W \subset \mathbb{R}^2$ (See [3], pp. 254-256) such that φ_2^* is an extension of φ_2 in W, and

$$P \in W, \quad \varphi_2^* \in C^1(W) \quad \text{and} \quad \varphi_2^*|_{\Omega \cap W} = \varphi_2, \quad \varphi_2^*|_{\partial\Omega \cap W} = 0. \tag{2.2}$$

Let $P = (x_0, y_0) \in \partial\Omega \cap N$. If $\dfrac{\partial \varphi_2}{\partial \nu}(P) \neq 0$, then $\pm|\nabla \varphi_2^*|(P) = \nabla \varphi_2^* \cdot \nu(P) = \dfrac{\partial \varphi_2^*}{\partial \nu}(P) = \dfrac{\partial \varphi_2}{\partial \nu}(P) \neq 0$ (Please see Appendix for the first equality). Without loss of generality, we assume $\dfrac{\partial \varphi_2^*}{\partial y}(P) \neq 0$. By Implicit Function Theorem, there exists a unique function $g : (x_0 - \epsilon, x_0 + \epsilon) \to \mathbb{R}$ (ϵ is small enough) such that $g \in C^1(x_0 - \epsilon, x_0 + \epsilon)$, $g(x_0) = y_0$ and for any $x \in (x_0 - \epsilon, x_0 + \epsilon)$ we have $\varphi_2^*(x, g(x)) = \varphi_2^*(x_0, y_0) = 0$. Which implies that $(x, g(x)) \in \partial\Omega$ by (2.3), hence $\varphi_2^*(x, y) \neq 0$ for $(x, y) \in \Omega \cap W$. Which contradict to $P \in N$. i.e. $\dfrac{\partial \varphi_2}{\partial \nu}(P) = \pm|\nabla \varphi_2^*|(P) = 0$.

Secondly, suppose $P \in N$, then, by the Hopf's Lemma, we have $\dfrac{\partial \varphi_2}{\partial \nu}(P) \neq 0$. $\square$

By the same way, we obtain the following Lemma 2:

Lemma 2 Suppose $\widetilde{P} \in \partial\widetilde{\Omega}$. Then $\dfrac{\partial \widetilde{\varphi}_2}{\partial \widetilde{\nu}}(\widetilde{P}) = 0$ if and only if $\widetilde{P} \in \widetilde{N}$, where $\dfrac{\partial \widetilde{\varphi}_2}{\partial \widetilde{\nu}}$ is the outnormal derivative of $\widetilde{\varphi}_2$ on the boundary.

Proof Let $\widetilde{P} \in \partial\widetilde{\Omega}$. Since $\widetilde{\varphi}_2$ is a smooth solution of equation (2.1) and $\widetilde{\Omega}$ is a smoothly bounded domain, there exists an open set $\widetilde{W} \subset \mathbb{R}^2$ such that $\widetilde{\varphi}_2^*$ is an extension of $\widetilde{\varphi}_2$ in $\widetilde{W}$, and

$$\widetilde{P} \in \widetilde{W}, \quad \widetilde{\varphi}_2^* \in C^1(\widetilde{W}) \quad \text{and} \quad \widetilde{\varphi}_2^*|_{\widetilde{\Omega} \cap \widetilde{W}} = \widetilde{\varphi}_2, \quad \widetilde{\varphi}_2^*|_{\partial\widetilde{\Omega} \cap \widetilde{W}} = 0. \tag{2.3}$$

Let $\widetilde{P} = (r_0, \theta_0) \in \partial\widetilde{\Omega} \cap \widetilde{N}$. If $\dfrac{\partial \widetilde{\varphi}_2}{\partial \widetilde{\nu}}(\widetilde{P}) \neq 0$, then $\pm|\nabla \widetilde{\varphi}_2^*|(\widetilde{P}) = \nabla \widetilde{\varphi}_2^* \cdot \widetilde{\nu}(\widetilde{P}) = \dfrac{\partial \widetilde{\varphi}_2^*}{\partial \widetilde{\nu}}(\widetilde{P}) = \dfrac{\partial \widetilde{\varphi}_2}{\partial \widetilde{\nu}}(\widetilde{P}) \neq 0$. Without loss of generality, we assume $\dfrac{\partial \widetilde{\varphi}_2^*}{\partial \theta}(\widetilde{P}) \neq 0$. By Implicit Function Theorem, there exists a unique function $g : (r_0 - \epsilon, r_0 + \epsilon) \to \mathbb{R}$

(ϵ is small enough) such that $g \in C^1(r_0 - \epsilon, r_0 + \epsilon)$, $g(r_0) = \theta_0$ and for any $r \in (r_0 - \epsilon, r_0 + \epsilon)$ we have $\widetilde{\varphi}_2^*(r, g(r)) = \widetilde{\varphi}_2^*(r_0, \theta_0) = 0$. Which implies that $(r, g(r)) \in \partial\widetilde{\Omega}$ by (2.3), hence $\widetilde{\varphi}_2^*(r, \theta) \neq 0$ for $(r, \theta) \in \widetilde{\Omega} \cap \widetilde{W}$. Which contradict to $\widetilde{P} \in \widetilde{N}$. i.e. $\dfrac{\partial \widetilde{\varphi}_2}{\partial \widetilde{\nu}}(\widetilde{P}) = \pm|\nabla \widetilde{\varphi}_2^*|(\widetilde{P}) = 0$.

Now, we show that if $\dfrac{\partial \widetilde{\varphi}_2}{\partial \widetilde{\nu}}(\widetilde{P}) = 0$ then $\widetilde{P} \in \widetilde{N}$.

Otherwise, there exist a neighborhood $U \subset \mathbb{R}^2$ of $\widetilde{P}$ such that $N \cap U = \emptyset$. Then either $\widetilde{\varphi}_2(r, \theta) > 0$ or $\widetilde{\varphi}_2(r, \theta) < 0$ for all $(r, \theta) \in U \cap \widetilde{\Omega}$, without loss of generality, we assume that $\widetilde{\varphi}_2(r, \theta) < 0$ for all $(r, \theta) \in U \cap \widetilde{\Omega}$. Since $\widetilde{\varphi}_2$ divides the domain $\widetilde{\Omega}$ into at exactly 2 subregions $\widetilde{\Omega}_1$ and $\widetilde{\Omega}_2$, we take $U \cap \widetilde{\Omega} \subset \widetilde{\Omega}_1$, then $\widetilde{\varphi}_2(r, \theta) < 0$ for all $(r, \theta) \in \widetilde{\Omega}_1$. Since $\Big(- \dfrac{\partial^2}{\partial r^2} - \dfrac{1}{r^2} \dfrac{\partial^2}{\partial \theta^2} - \dfrac{1}{r} \dfrac{\partial}{\partial r} \Big)(\widetilde{\varphi}_2) = \lambda_2 \widetilde{\varphi}_2 \leq 0$ in $\widetilde{\Omega}_1$, and $\widetilde{\varphi}_2(\widetilde{P}) = 0 > \widetilde{\varphi}_2(r, \theta)$ for all $(r, \theta) \in \widetilde{\Omega}_1$, we obtain $\dfrac{\partial \widetilde{\varphi}_2}{\partial \widetilde{\nu}}(\widetilde{P}) > 0$ by Hopf's Lemma (See [3] pp. 330-332). Which implies the contradiction. $\square$

Now we consider the equation:

$$\begin{cases} \dfrac{\partial^2 \widetilde{\varphi}_2}{\partial r^2} + \dfrac{1}{r^2} \dfrac{\partial^2 \widetilde{\varphi}_2}{\partial \theta^2} + \dfrac{1}{r} \dfrac{\partial \widetilde{\varphi}_2}{\partial r} + \lambda_2 \widetilde{\varphi}_2 = 0 & \text{in } \widetilde{\Omega} \\ \widetilde{\varphi}_2 = 0 & \text{on } \partial\widetilde{\Omega}, \end{cases} \tag{2.4}$$

where $\widetilde{\Omega}$ is a smoothly bounded domain, $\widetilde{\Omega} \subset \{(r, \theta) \in \mathbb{R}^2;\ 0 < r_0 \leq r \leq R_0\}$ (r_0, R_0 are given constants), $\widetilde{\Omega}$ is strictly convex in θ and $\widetilde{\Omega}$ is symmetric with respect to r-axis. (We note that the domain $\widetilde{\Omega}$ is strictly convex in θ if every line parallel to the θ-axis which intersect $\widetilde{\Omega}$, cut $\partial\Omega$ in at most two points.)

Then we obtain the following Lemma 3:

Lemma 3 The nodal line $\widetilde{N}$ of a second eigenfunction $\widetilde{\varphi}_2$ divides the domain $\widetilde{\Omega}$ by intersecting its boundary at exactly two points.

Proof The proof of this lemma is similar to Theorem 2.2 in [8] and Theorem I in [11].

If $\widetilde{\varphi}_2$ is odd in θ (i.e. $\widetilde{\varphi}_2(r, -\theta) = -\widetilde{\varphi}_2(r, \theta)$), the nodal line $\widetilde{N}$ is just the r-axis. And Lemma 3 is obviously true. Suppose $\widetilde{\varphi}_2$ is even in r (i.e. $\widetilde{\varphi}_2(r, -\theta) = \widetilde{\varphi}_2(r, \theta)$). Assume that Lemma 3 is false. Then for $\widetilde{P} \in \partial\widetilde{\Omega}$, $(\dfrac{\partial \widetilde{\varphi}}{\partial \theta})(\widetilde{P}) \neq 0$, except the tangent of $\partial\Omega$ at $\widetilde{P}$ is the θ-direction. Without loss of generality, we may assume that $(\dfrac{\partial \widetilde{\varphi}}{\partial \theta})(\widetilde{P}) \geq 0$ for $\widetilde{P} \in \partial\widetilde{\Omega} \cap \{(r, \theta);\ \theta \leq 0\}$. Set $(\widetilde{\Omega})^- = \mathrm{Cl}(\widetilde{\Omega}) \cap \{(r, \theta);\ \theta \leq 0\}$, here and throughout the paper $\mathrm{Cl}(\widetilde{\Omega})$ represent the closure of $\widetilde{\Omega}$. $\dfrac{\partial \widetilde{\varphi}}{\partial \theta}$ must change sign in $(\widetilde{\Omega})^-$. Otherwise $\widetilde{\varphi}_2 \geq 0$ in $(\widetilde{\Omega})^-$. by evenness, $\widetilde{\varphi}_2 \geq 0$ in $\mathrm{Cl}(\widetilde{\Omega})$, which leads to a contradiction. Hence the nodal line $\{(r,\theta) \in (\widetilde{\Omega})^-; \ \frac{\partial \widetilde{\varphi}}{\partial \theta}(r,\theta) = 0\}$ encloses a subregion $(\widetilde{\Omega})^-_*$ of $\{(r,\theta) \in (\widetilde{\Omega})^-; \ \frac{\partial \widetilde{\varphi}}{\partial \theta}(r,\theta) < 0\}$. Let $\widetilde{\Omega}_* = \{(r,\theta) \in \widetilde{\Omega}; \text{ either } (r,\theta) \in (\widetilde{\Omega})^-_* \text{ or } (r,-\theta) \in (\widetilde{\Omega})^-_*\}$. Then $\frac{\partial \widetilde{\varphi}_2}{\partial \theta}$ satisfies

$$\begin{aligned}&\Big(\frac{\partial^2}{\partial r^2} + \frac{1}{r^2}\frac{\partial^2}{\partial \theta^2} + \frac{1}{r}\frac{\partial}{\partial r} + \lambda_2\Big)(\frac{\partial \widetilde{\varphi}_2}{\partial \theta}) \\ &= \frac{\partial}{\partial \theta}\Big[\Big(\frac{\partial^2}{\partial r^2} + \frac{1}{r^2}\frac{\partial^2}{\partial \theta^2} + \frac{1}{r}\frac{\partial}{\partial r} + \lambda_2\Big)(\widetilde{\varphi}_2)\Big] = 0 \quad \text{in } \widetilde{\Omega}_*,\end{aligned} \tag{2.5}$$

and

$$\frac{\partial \widetilde{\varphi}_2}{\partial \theta} = 0 \quad \text{on } \partial\widetilde{\Omega}_*. \tag{2.6}$$

By (2.5) and (2.6) we obtain that $\widehat{\varphi}_2 := \frac{\partial \widetilde{\varphi}_2}{\partial \theta} \circ \rho$ satisfies

$$\begin{cases}\Delta \widehat{\varphi}_2 + \lambda_2 \widehat{\varphi}_2 = 0 & \text{in } \Omega_* := \rho^{-1}(\widetilde{\Omega}_*) \\ \widehat{\varphi}_2 = 0 & \text{on } \partial\Omega_*,\end{cases}$$

Since $\frac{\partial \widetilde{\varphi}_2}{\partial \theta}(r,-\theta) = -\frac{\partial \widetilde{\varphi}_2}{\partial \theta}(r,\theta)$, $\frac{\partial \widetilde{\varphi}_2}{\partial \theta}$ must change sign, Then $\widehat{\varphi}_2$ is also change sign. Hence $\lambda_2 \geq \lambda_2(\Omega_*)$, where $\lambda_2(\Omega_*)$ is the second eigenvalue of Laplacian in Ω_*. But $\Omega_* \subset \Omega$ (since $\widetilde{\Omega}_* \subset \widetilde{\Omega}$), and by monotony principle we get

$$\lambda_2 \geq \lambda_2(\Omega_*) > \lambda_2,$$

which is a contradiction.

In general a second eigenfunction $\widetilde{\varphi}_2$ can be written as $\widetilde{\varphi}_2 = \phi_1 + \phi_2$, where $\phi_1(r,\theta) = \frac{\widetilde{\varphi}_2(r,\theta) - \widetilde{\varphi}_2(r,-\theta)}{2}$ is odd in θ, $\phi_2(r,\theta) = \frac{\widetilde{\varphi}_2(r,\theta) + \widetilde{\varphi}_2(r,-\theta)}{2}$ is even in θ and both are second eigenfunctions. By the above proof, we know that there exist two points $\widetilde{P} = (r_0,\theta_0)$ and $\widetilde{Q} = (r_0,-\theta_0) \in \partial\widetilde{\Omega}$, where $\theta_0 \neq 0$ such that $(\frac{\partial \phi_2}{\partial \widetilde{\nu}})(\widetilde{P}) = (\frac{\partial \phi_2}{\partial \widetilde{\nu}})(\widetilde{Q}) = 0$. Since $\frac{\partial \phi_1}{\partial \widetilde{\nu}}$ is odd on $\partial\widetilde{\Omega}$, we assume $(\frac{\partial \phi_1}{\partial \widetilde{\nu}})(\widetilde{P}) > 0$ and $(\frac{\partial \phi_1}{\partial \widetilde{\nu}})(\widetilde{Q}) < 0$. Hence $(\frac{\partial \widetilde{\varphi}_2}{\partial \widetilde{\nu}})(\widetilde{P}) > 0$ and $(\frac{\partial \widetilde{\varphi}_2}{\partial \widetilde{\nu}})(\widetilde{Q}) < 0$. It implies that there exist two points $\widetilde{P}^*$ and $\widetilde{Q}^*$ on $\partial\widetilde{\Omega}$ such that

$$(\frac{\partial \phi_2}{\partial \widetilde{\nu}})(\widetilde{P}^*) = (\frac{\partial \phi_2}{\partial \widetilde{\nu}})(\widetilde{Q}^*) = 0,$$

and the Lemma 3 for this case follows. $\square$

Since $\rho : \Omega \to \widetilde{\Omega},\ (x, y) \mapsto (r, \theta)$ is a diffeomorphism defined by $(x, y) = \rho^{-1}(r, \theta)$ with $x = r\cos\theta, y = r\sin\theta$, we get the following theorem:

Theorem The nodal line of a second eigenfunction of Laplacian divides the domain Ω by intersecting its boundary at exactly two points if the domain $\rho(\Omega)$ is strictly convex in θ and symmetric with respect to the r-axis.

Remark At the end of this paper, we give a figure, which displays that the diffeomorphism $\rho(\Omega)$ of some concave domain Ω satisfies the conditions of Theorem.

In the following figure, Ω in figure (A) is the domain formed by the curves C_1, C_2, C_3, C_4, where $C_1 = \{(x, y) \in \mathbb{R}^2;\ \frac{x^2}{6^2} + \frac{y^2}{3^2} = 1 \text{ and } x \geq 0\}$, $C_2 = \{(x, y) \in \mathbb{R}^2;\ \frac{x^2}{1^2} + \frac{y^2}{2^2} = 1 \text{ and } x \geq 0\}$, $C_3 = \{(x, y) \in \mathbb{R}^2;\ x^2 + (y - \frac{5}{2})^2 = (\frac{1}{2})^2 \text{ and } x \leq 0\}$, $C_4 = \{(x, y) \in \mathbb{R}^2;\ x^2 + (y + \frac{5}{2})^2 = (\frac{1}{2})^2 \text{ and } x \leq 0\}$. $\rho(\Omega)$ in figure (B) is the diffeomorphism of Ω. Obviously, Ω is a concave domain and $\rho(\Omega)$ is a convex domain.



References

[1] G. Alessandrini, Nodal lines of eigenfunctions of the fixed membrane problem in general convex domains, Comment. Math. helvetici 69 (1994), 142-154.

[2] R. Courant and D. Hilbert, Methods of mathemathical physis (Vol. 1), New York, 1953.

[3] L. C. Evans, Partial Differential Equations (Grduate Studies in Mathematics 19), AMS providence, Rhode Island, 1997.

[4] D. Grieser and D. Jerison, Asymptotics of the first nodal line of a convex domain, Invent. Math. 125 (1996), 197-219.

[5] M. Hoffmann-Ostenhof, T. M. Hoffmann-Ostenhof and N. Nadirashvili, The nodal line of the second eigenfunction of the Laplacian in $\mathbb{R}^2$ can be closed, Duke Math. J., Vol. 90, no. 3 (1997), 631-640.

[6] D. Jerison, The first nodal line of a convex planar domain, Internat. Math. Res. Notices, No. 1 (1991), 1-5; Duke Math. J. 62, 1991.

[7] D. Jerison, The diameter of the first nodal line of a convex domain, Annal of Math. 141 (1995), 1-33.

[8] Chang-Shou Lin, On the second eigenfunctions of the Laplacian in $\mathbb{R}^2$, commun. Math. Phys. 111 (1987), 161-166.

[9] A. D. Melas, On the nodal line of the second eigenvalues of the Laplacian in $\mathbb{R}^2$, J. Diff. Geom. 35 (1992), 255-263.

[10] L. Payne, Isoperimtric inequalities and their applications, Siam Review 9 (1967), 453-488.

[11] L. Payne, On two conjectures in the fixed membrance eigenvalue problem, J. Appl. Math. Phys. (ZAMP) 24 (1973), 720-729.

[12] S. -T. Yau, Problem section in 'Seminar on differential geometry', Ann. Math. Studies 102 (1982), 669-706.

Appendix

Claim Let $\Omega \subset \mathbb{R}^n$ $(n \in \mathbb{Z}^+)$ be a smooth domain and $P \in \partial\Omega$. Let W be any open set containing P. Then for any C^1 function $\varphi : W \to \mathbb{R}$ with $\varphi|_{\partial\Omega\cap W} = 0$ and $(\nabla\varphi)(P) \neq 0$ we have $\nu(P) = \pm\dfrac{\nabla\varphi}{|\nabla\varphi|}(P)$, where $\nu(P)$ is the outer normal vector of $\partial\Omega$ at P.

Proof of Claim Since $\varphi|_{\partial\Omega\cap W} = 0$ and $(\nabla\varphi)(P) \neq 0$, there exists a neighborhood of P, without loss of generality, we assume the neighborhood is W, such that $\{x \in W;\ \varphi(x) = 0\} = \partial\Omega \cap W$, i.e. $\partial\Omega \cap W$ is a regular surface. Let $v \in \mathbb{R}^n$ be any normal tangent vector of $\partial\Omega$ at P, there exists a curve $\gamma : (-\epsilon, \epsilon) \to \mathbb{R}^n$ such that $\gamma(-\epsilon, \epsilon) \subset \partial\Omega$ and $\gamma(0) = P, \gamma'(0) = v$. Then $\varphi(\gamma(t)) = 0$ and $0 = \dfrac{d}{dt}|_{t=0}\varphi(\gamma(t)) = (\nabla\varphi)|_{\gamma(0)} \cdot \gamma'(0) = (\nabla\varphi)(P) \cdot v$. Hence $(\nabla\varphi)(P) = [\pm|\nabla\varphi|(P)]\nu(P)$. We have prove the Claim.